

TABLE 1. Total Numbers and Relative Species Composition of Anadromous Salmonids Trapped in the South Fork Trinity River Near Sandy Bar, 1984 through 1990 (nd = no data).

Sampling dates	1984		1985		1986		1987	
	No.	%	No.	%	No.	%	No.	%
09/17-11/02	46		61		98		69	
No. sampling days								
Chinook salmon								
Fall-run	73	55.7	176	35.6	264	39.0	455	63.6
Spring-run	nd	-	nd	-	13	1.9	nd	
Coho salmon	3	2.3	109	22.0	12	1.8	17	2.4
Steelhead	55	42.0	207	41.8	387	57.2	243	33.9
Chum salmon	0	0.0	3	0.6	1	0.1	1	0.1
Brown trout	0	0.0	0	0.0	0	0.0	0	0.0
TOTALS	131		495		677		716	

Sampling dates	1988		1989		1990	
	No.	%	No.	%	No.	%
09/22-11/13	52		39		169	
No. sampling days						
Chinook salmon						
Fall-run	368	61.5	52	57.8	223	48.3
Spring-run	nd	-	nd	-	nd	-
Coho salmon	3	0.5	1	1.1	61	13.2
Steelhead	227	38.0	37	41.1	176	38.1
Chum salmon	0	0.0	0	0.0	1	0.2
Brown trout	0	0.0	0	0.0	1	0.2
TOTALS	598		90		462	